

Institute of Geometry

## Advanced Topics in Discrete Mathematics Seminar

Fr 28.06.2024, 11:00 Uhr

Seminarraum 2, Kopernikusgasse 24/IV.

# The excedance quotient of the Bruhat order, Quasisymmetric Varieties and Temperley-Lieb algebras

NANTEL BERGERON  
(York University, Canada)

Let  $R_n = \mathbb{Q}[x_1, x_2, \dots, x_n]$  be the ring of polynomials in  $n$  variables and consider the ideal  $\langle \text{QSym}_n^+ \rangle \subseteq R_n$  generated by quasisymmetric polynomials without constant term. It was shown by J. C. Aval, F. Bergeron and N. Bergeron that  $\dim(R_n / \langle \text{QSym}_n^+ \rangle) = C_n$  the  $n$ th Catalan number. In the present work, we explain this phenomenon by defining a set of permutations  $\text{QSV}_n$  with the following properties: first,  $\text{QSV}_n$  is a basis of the Temperley-Lieb algebra  $\text{TL}_n(2)$ , and second, when considering  $\text{QSV}_n$  as a collection of points in  $\mathbb{Q}^n$ , the top-degree homogeneous component of the vanishing ideal  $\mathbf{I}(\text{QSV}_n)$  is  $\langle \text{QSym}_n^+ \rangle$ .

Our construction has a few byproducts which are independently noteworthy. We define an equivalence relation  $\sim$  on the symmetric group  $S_n$  using weak excedances and show that its equivalence classes are naturally indexed by noncrossing partitions. Each equivalence class is an interval in the Bruhat order between an element of  $\text{QSV}_n$  and a 321-avoiding permutation. Furthermore, the Bruhat order induces a well-defined order on  $S_n / \sim$ . Finally, we show that any section of the quotient  $S_n / \sim$  gives an (often novel) basis for  $\text{TL}_n(2)$ .

This talk is based on joint work with Lucas Gagnon.

Cesar Ceballos